Orzeyful™
oveporexton tablets (IV)



LET'S TALK ABOUT

narcolepsy type 1
(narcolepsy with cataplexy)

Narcolepsy type 1 (NT1) can be hard to explain—especially when symptoms can change daily. The more clearly you describe what you're experiencing, the easier it is for your healthcare provider to help find a management plan that's right for you.

Actor portrayal.

A guide to help start the conversation with your healthcare provider

What is ORZEYFUL?

ORZEYFUL™ is a prescription medicine used in adults to treat narcolepsy type 1 (narcolepsy with cataplexy).

It is not known if ORZEYFUL is safe and effective in children under 18 years of age.

IMPORTANT SAFETY INFORMATION

Do not take ORZEYFUL if you take certain medicines that are strong inhibitors of an enzyme called CYP3A, since they may increase your risk of side effects. Ask your healthcare provider if you are not sure.

Please see additional Important Safety Information throughout and click for full Prescribing Information and talk with your healthcare provider.

Sharing all NT1 symptoms with your healthcare provider can help them understand what you are going through

1. **What symptoms are you dealing with? How often do they occur?**
   (Please fill out the chart below according to your symptoms)

Symptoms	How many days a week?	When does this typically happen? (e.g., time of day, during certain activities, or in specific situations)
Trouble staying awake during the day (excessive daytime sleepiness)		
A sudden loss of muscle control (cataplexy)		
Inability to move or speak when falling asleep or waking up (sleep paralysis)		
Falling asleep quickly, but waking up often (disrupted nighttime sleep)		
Vivid, dreamlike experiences when falling asleep or waking (hallucinations)		

Other _____

_____

_____

Sharing all NT1 symptoms with your healthcare provider can help them understand what you are going through

2. **What do your symptoms feel like?**

3. **What about NT1 can feel most challenging for you?**

4. **How does NT1 impact you?**

- I'm getting by, but things could be better. (Please describe)
- I'm struggling and want to discuss managing my NT1. (Please describe)
- I'm greatly impacted by NT1. (Please describe)

5. **In the past, what ways have you tried to manage your symptoms?**

- Behavioral coping strategies (Please describe)
- Medication(s) (Please list)

6. **What are the symptoms you would like to better manage?**

A few conversation starters to get the discussion going

In your own words, describe your personal experiences so your healthcare provider can better understand what you're going through.

Use these sentence starters (fill in the blanks):

"My sleepiness can sometimes _____

_____ ."

"The times when I feel the most tired are _____

_____ ."

"In a typical week, I can have planned naps _____ times and unplanned naps _____ times."

"Sometimes I feel like _____

_____ ."

"I'm concerned about _____ because _____

_____ ."

Evaluate your level of daytime sleepiness and bring the results to your healthcare provider

The Epworth Sleepiness Scale (ESS) is a short questionnaire that helps measure how likely you are to doze off in everyday situations.

Click here to **take the ESS**

"I really want to know my patients' day-to-day. Not only what's going well, but also what they're struggling with.

—From real healthcare providers

"Everyone is different. So, I share all my experiences with my doctor so they can help me better manage my NT1.

—A real NT1 patient



Actor portrayal.

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of ORZEYFUL?

- **ORZEYFUL may cause serious side effects, including:**
  - **Trouble sleeping (insomnia).** Tell your healthcare provider if you develop insomnia, especially if it does not go away after 7 days, while taking ORZEYFUL. They may decrease your dose or tell you to stop taking ORZEYFUL.
  - **Urinary frequency and urgency.** Tell your healthcare provider if you urinate more often or have sudden urges to urinate. If you have a history of overactive bladder, your healthcare provider may monitor you for worsening urinary symptoms.
  - **Increased level of an enzyme in your blood called creatine phosphokinase (CPK).** Increased CPK may be a sign of muscle damage. Tell your healthcare provider right away if you have unexplained muscle pain, weakness, or dark urine, especially after vigorous physical activity, or if you take other medicines that may cause muscle problems.
- **Most common side effects of ORZEYFUL include** trouble sleeping (insomnia), increased urinary frequency, urinary urgency, and increased saliva.

  These are not all the possible side effects of ORZEYFUL. Call your healthcare provider for medical advice about side effects.

Before taking ORZEYFUL, tell your healthcare provider about all your medical conditions, including if you:

- have a history of overactive bladder
- have a history of drug or alcohol abuse or addiction
- have severe liver or kidney problems
- are pregnant or plan to become pregnant. It is not known if ORZEYFUL will harm your unborn baby.
  - There is a pregnancy registry for women who are exposed to ORZEYFUL during pregnancy. The registry collects information about the health of you and your baby. To enroll or get information from the registry, call 1-877-371-0309.
- are breastfeeding or plan to breastfeed. Talk to your healthcare provider about the best way to feed your baby if you take ORZEYFUL.

Tell your healthcare provider about all the medicines you take, including prescription medicines and over-the-counter medicines, vitamins, and herbal supplements. Some medicines may affect how ORZEYFUL works or increase your risk of side effects. Do not start, stop, or change any other medicine while taking ORZEYFUL without talking to your healthcare provider.

Please see additional Important Safety Information throughout and click for full Prescribing Information and talk with your healthcare provider.

Orzeyful™
oveporexton tablets Ⓒ

Orzeyful™
oveporexton tablets (IV)

INTRODUCING

A new treatment option for adults with narcolepsy type 1 (narcolepsy with cataplexy)



Actor portrayal.

Only ORZEYFUL targets the underlying loss of orexin in NT1 by activating orexin receptors in the brain to restore orexin signaling.

Talk to your healthcare provider to see if ORZEYFUL may be right for you.

Individual results may vary.

Learn more at **orzeyful.com**

IMPORTANT SAFETY INFORMATION (cont'd)

ORZEYFUL is a federally controlled substance (CIV) because it can be abused or lead to dependence. Tell your healthcare provider if you have ever abused or been dependent on alcohol, prescription medicines, or street drugs. Keep your ORZEYFUL in a safe place and never give it to anyone else.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout and click for full Prescribing Information and talk with your healthcare provider.

ORZEYFUL is available as 0.5 mg, 1 mg, and 2 mg tablets.

Takeda

©2026 Takeda Pharmaceuticals U.S.A., Inc. 1-877-TAKEDA-7 (1-877-825-3327). All rights reserved. ORZEYFUL and The ORZEYFUL Logo are trademarks of Takeda Pharmaceuticals International AG. Takeda and Takeda are registered trademarks of Takeda Pharmaceutical Company Limited. US-OVE-0187v1.0 11/26